

## PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/108879>

Please be advised that this information was generated on 2017-12-06 and may be subject to change.

*Critical Review***Nucleus Accumbens and Dopamine-Mediated Turning Behavior of the Rat: Role of Accumbal Non-dopaminergic Receptors**Hiroko Ikeda<sup>1,2,\*</sup>, Junzo Kamei<sup>2</sup>, Noriaki Koshikawa<sup>1,3</sup>, and Alexander R. Cools<sup>4</sup><sup>1</sup>Department of Pharmacology, Nihon University School of Dentistry,  
1-8-13 Kanda-Surugadai, Chiyoda-ku, Tokyo 101-8310, Japan<sup>2</sup>Department of Pathophysiology and Therapeutics, Hoshi University School of Pharmacy and Pharmaceutical Sciences,  
2-4-41 Ebara, Shinagawa-ku, Tokyo 142-8501, Japan<sup>3</sup>Division of Oral and Craniomaxillofacial Research, Dental Research Center, Nihon University School of Dentistry,  
1-8-13 Kanda-Surugadai, Chiyoda-ku, Tokyo 101-8310, Japan<sup>4</sup>Radboud University Medical Centre Nijmegen, Donders Institute for Brain,  
Cognition and Behavior: Psychoneuropharmacology, P.O. Box 9101, 6500 HB Nijmegen, The Netherlands

Received March 26, 2012; Accepted August 20, 2012

**Abstract.** Accumbal dopamine plays an important role in physiological responses and diseases such as schizophrenia, Parkinson's disease, and depression. Since the nucleus accumbens contains different neurotransmitters, it is important to know how they interact with dopaminergic function: this is because modifying accumbal dopamine has far-reaching consequences for the treatment of diseases in which accumbal dopamine is involved. This review provides a summary of these interactions, and our current knowledge about them are as follows: A) AMPA receptors are required for dopamine-dependent behavior and *vice versa*; NMDA receptors modulate the activity at the level of AMPA and/or dopamine D<sub>1</sub> receptors. B) GABA<sub>A</sub>, but not GABA<sub>B</sub>, receptors inhibit dopamine-dependent behavior. C) Nicotinic receptors are required for dopamine-dependent behavior, whereas muscarinic receptors inhibit dopamine-dependent behavior. D)  $\alpha$ -Adrenoceptors inhibit dopamine-dependent behavior in contrast to  $\beta$ -adrenoceptors, which potentiate this behavior. E)  $\mu$ - and  $\delta$ <sub>2</sub>-opioid receptors elicit behavior that requires an intact dopaminergic function and  $\delta$ <sub>2</sub>-opioid receptors modulate dopamine-dependent behavior. F) Orexin 2 receptors play an important, modifying role in dopamine-dependent behavior. G) Somatostatin receptors potentiate dopamine-dependent behavior. It is suggested that modulation of the above-mentioned non-dopaminergic receptors provide new tools to control physiological functions as well as diseases mediated by accumbal dopamine.

**Keywords:** dopamine, nucleus accumbens, turning behavior

**1. Introduction**

Dopamine in the central nervous system plays an important role not only in behavior and physiological responses, but also in many diseases (review: 1; depression: 2; drug addiction: 3 – 5; attention deficit hyperactivity disorder (ADHD): 6, 7; schizophrenia: 8, 9; Parkinson's

disease, review: 10). The nucleus accumbens (Nacc), which is a terminal area of the mesolimbic dopaminergic neurons, is known to be one of the main structures controlling physiological responses, behavior, and diseases (1). The Nacc is a heterogeneous structure and is divided anatomically into at least two structures, that is, the shell and the core (11 – 14). There are many differences between these two structures. For instance, the dopamine plexus is richer in the shell than in the core (13), and the concentration of dopamine is higher in the shell than in the core (12). Furthermore, the dopamine D<sub>1</sub> binding in the rostral areas of the Nacc appears to be higher in the shell than in the core, whereas the dopamine D<sub>2</sub> binding

\*Corresponding author (affiliation #2). h-ikeda@hoshi.ac.jp  
Published online in J-STAGE on October 10, 2012 (in advance)  
doi: 10.1254/jphs.12R02CR

appears to be lower in the shell than in the core (11). Thus, it is not remarkable that dopamine functions vary across the core and the shell (15, 16). The shell is thought to play a pivotal role in dopamine-mediated functions because it contains the largest amount of dopaminergic terminals and the highest concentration of dopamine (12, 13).

Unilateral activation of the Nacc has been shown to elicit two characteristic types of turning behavior, namely circling and pivoting (16, 17). Circling is marked by normal hindlimb stepping, normal forelimb stepping, turns of large diameter ( $> 30$  cm), and running, whereas pivoting is marked by abnormal hindlimb stepping, turns of small diameter ( $< 20$  cm), and spinning around one hindlimb (17). More specifically, we have provided pharmacological evidence that unilateral stimulation of dopamine  $D_1/D_2$  receptors in the Nacc shell, but not the Nacc core, elicits contraversive pivoting, whereas unilateral stimulation of acetylcholine receptors in the Nacc shell elicits contraversive circling (15, 16). Since the contraversive pivoting induced by unilateral stimulation of dopamine  $D_1/D_2$  receptors in the Nacc shell is a characteristic feature of the function of accumbal dopamine in the Nacc shell and the contraversive circling induced by unilateral stimulation of acetylcholine receptors in the Nacc shell is a characteristic feature of the function of accumbal acetylcholine in the Nacc shell, these behaviors are very useful tools for studies on analysis of interactions between various accumbal, neuroactive compounds as well as for studies on analysis of the involved output stations of the Nacc. We have provided evidence that the shell-specific, dopaminergic pivoting is mediated via a substrate that differs from that involved in the shell-specific, cholinergic circling (16, 18–21).

The Nacc receives not only dopaminergic projections from the ventral tegmental area (VTA) and the substantia nigra pars compacta (SNc), but also many other neuronal projections releasing other neurotransmitters and neuropeptides (see below). It is important to know to what extent these neuroactive compounds direct the function of accumbal dopamine: this is because accumbal dopamine has important therapeutic consequences for the treatment of diseases in which accumbal dopamine is involved. This review provides a summary of these interactions. Since a single neurotransmitter or neuropeptide can act at distinct receptors, this review focuses on the involved receptors.

## 2. Interaction with dopamine in the Nacc

Because the Nacc receives a variety of neuronal projections, each releasing their own neurotransmitter and/or neuropeptide, it is not surprising that mutual interac-

tions between dopamine and neurotransmitters such as glutamate,  $\gamma$ -aminobutyric acid (GABA), acetylcholine, noradrenaline, serotonin, and neuropeptides such as enkephalin, orexin, and somatostatin occur in the Nacc (see below). These interactions as revealed by studying how compounds that selectively interact with one of the various accumbal, non-dopaminergic receptors influence the shell-specific, dopaminergic pivoting, are discussed below. Recently, heterodimerization of dopamine receptors has been shown for both the  $D_1/D_5$ - and  $D_2/D_3/D_4$ -receptor families, which couple positively and negatively, respectively, to adenylyl cyclase (for reviews: 22, 23; see also: 24). Notably, heterodimers are formed by  $D_1$  and adenosine  $A_1$  receptors,  $D_2$  or  $D_3$  and adenosine  $A_2$  receptors,  $D_2$  and somatostatin  $SST_5$  receptors, and  $D_2$  and cannabinoid  $CB_1$  receptors. Further,  $D_1$ ,  $D_2$ , and  $D_3$  receptors physically assemble into functional  $D_1/D_2$ ,  $D_1/D_3$ , and  $D_2/D_3$  heterodimers possessing binding and coupling profiles distinct from the respective monomers. By using different experimental approaches various groups have, in fact, demonstrated the existence of dopamine heterodimers in both transfected cell systems and the striatum with peculiar pharmacological, signaling, and functional properties. The putative role of these heteromers in the physiological regulation of accumbal function is yet unknown. For that reason, the role of these heteromers will not be discussed, although future research may show that these have a functional and therapeutic significance.

### 2.1. Glutamate


The Nacc receives glutamatergic inputs from several brain areas including the prefrontal cortex, amygdala, and hippocampus (25–27). Anatomical evidence indicates that there are interactions between dopaminergic and glutamatergic inputs in the Nacc (28, 29). The ionotropic glutamate receptors are classified into three types, that is, *N*-methyl-D-aspartic acid (NMDA),  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolpropionic acid (AMPA), and kainate receptors (30). Glutamatergic agonists have been found to enhance dopamine release in the Nacc, possibly via non-NMDA receptors (31), whereas accumbal dopamine has been found to increase extracellular glutamate levels in the Nacc (32). In behavioral studies, agonists of the glutamatergic ionotropic receptors, AMPA and NMDA receptors, have been found to stimulate locomotor activity when injected into the Nacc, and drugs that inhibit dopaminergic neurotransmission have been found to inhibit these effects (33, 34). Thus, both accumbal glutamate and accumbal dopamine regulate motor function.

To analyze in more detail the functional interaction between glutamate and dopamine in the Nacc shell, we have examined the role of glutamate receptors in eliciting

the shell-specific, dopaminergic pivoting. Unilateral injection of AMPA into the Nacc shell elicited contraversive pivoting and this effect was inhibited by the AMPA-receptor antagonist NBQX (1,2,3,4-tetrahydro-6-nitro-2,3-dioxo-benzo[*f*]quinoxaline-7-sulfonamide), indicating that unilateral stimulation of AMPA receptors in the Nacc shell elicits contraversive pivoting, which pattern is identical to that induced by unilateral stimulation of dopamine receptors in the Nacc shell (35). The contraversive pivoting induced by stimulation of AMPA receptors in the Nacc shell was inhibited by the dopamine D<sub>1</sub>/D<sub>2</sub>-receptor antagonist *cis*-(*Z*)-flupenthixol injected into the same area (35), indicating that the contraversive pivoting elicited by the stimulation of AMPA receptors in the Nacc shell requires intact dopaminergic neurotransmission in this brain region. Since AMPA-receptor stimulation has been found to enhance extracellular levels of dopamine in the Nacc (31), it has been suggested that stimulation of presynaptic AMPA receptors on dopaminergic terminals elicits contraversive pivoting (31, 35). In this context, it has to be stated that the stimulatory effects of AMPA receptors on dopamine release in the Nacc are not necessarily due to direct actions of this excitatory amino acid on dopaminergic terminals: indirect actions through intrinsic neurons or feedback loops cannot be excluded (36).

Stimulation or inhibition of NMDA receptors in the Nacc shell does not elicit turning behavior (35). The notion that NMDA receptors are not involved in this respect fits in with the finding that there is no functional interaction between accumbal NMDA receptors and accumbal dopamine receptors. In fact, evidence is available that local application of NMDA does not affect dopamine release from the Nacc (31). However, the NMDA-receptor antagonist MK-801 [(+)-5-methyl-10,11-dihydro-5*H*-dibenzo[*a,b*]cyclohepten-5,10-imine] has been found to inhibit dose-dependently the shell-specific, AMPA-induced pivoting when co-administered with AMPA into the Nacc shell (35), suggesting that NMDA receptors in the Nacc shell play a modulatory role in pivoting induced by unilateral injection of AMPA into the Nacc shell.

Contraversive pivoting induced by unilateral injection of a mixture of the dopamine D<sub>1</sub>-receptor agonist SKF 38393 and D<sub>2</sub>-receptor agonist quinpirole in the Nacc shell is inhibited by both MK-801 (Fig. 1A) and NBQX (Fig. 1B) injected into the same region (35). Thus, it has been concluded that the shell-specific, dopaminergic pivoting requires intact glutamatergic neurotransmission in the same region (35). This conclusion complements the earlier reported finding that locomotor activity produced by cocaine or dopamine D<sub>1</sub>/D<sub>2</sub>-receptor agonists are antagonized by intra-accumbens administration of AMPA- or NMDA-receptor antagonists (37, 38).



**Fig. 1.** Inhibition of contraversive pivoting induced by a mixture of SKF 38393 and quinpirole injected into the nucleus accumbens shell by intra-shell injections of MK-801 (A), NBQX (B), muscimol (C), mecamylamine (D), and phenylephrine (E) or by intraperitoneal injection of NTB (F). MK-801 (100 and 500 ng), NBQX (1 and 10 ng), muscimol (5 and 10 ng), mecamylamine (5 and 10 µg), and phenylephrine (5 and 10 µg) were injected into the nucleus accumbens shell as a cocktail with a mixture of SKF 38393 (5 µg) and quinpirole (10 µg). NTB (1 and 3 mg/kg) were injected intraperitoneally 30 min before the injection of a mixture of SKF 38393 (5 µg) and quinpirole (10 µg) into the nucleus accumbens shell. The contraversive pivoting was measured soon after the injection of a mixture of SKF 38393 and quinpirole into the nucleus accumbens shell. \* $P < 0.05$ , \*\* $P < 0.01$  vs. SKF 38393 + quinpirole group.

It has been suggested that stimulation of presynaptic dopaminergic receptors on glutamatergic terminals elicits contraversive pivoting identical to that elicited by stimulation of AMPA receptors in the Nacc shell (32, 35) because accumbal dopamine is known to increase the release of accumbal glutamate (32).

Recently, it has been reported that NMDA receptors

are localized on the neurons that contain dopamine D<sub>1</sub> receptors in the Nacc shell (39). Moreover, stimulation of dopamine D<sub>1</sub> receptors in the Nacc shell phosphorylates AMPA receptors, thereby strengthening the function of AMPA receptors (40). Therefore, an alternative explanation concerning the above-mentioned results is that NMDA and AMPA receptors postsynaptically modulate the dopamine function, possibly via D<sub>1</sub> receptors.

The above-mentioned results indicate that 1) accumbal AMPA receptors direct the shell-specific, dopaminergic pivoting, 2) accumbal, dopamine D<sub>1</sub>/D<sub>2</sub> receptors direct shell-specific, AMPA-mediated pivoting, and 3) accumbal NMDA receptors at best modulate the activity at the level of the involved accumbal AMPA and/or dopamine D<sub>1</sub> receptors.

## 2.2. GABA

The Nacc contains two types of GABA receptors, namely GABA<sub>A</sub> and GABA<sub>B</sub> receptors (41). GABA<sub>A</sub> receptors have been found to inhibit accumbal dopamine release, although GABA<sub>B</sub> receptors are not devoid of a modulatory role in this respect (42).

There is a close relationship between GABAergic and dopaminergic systems in the Nacc in terms of regulating locomotor behavior. For example, GABA<sub>A</sub>-receptor agonists reduce the hyperlocomotion induced by the dopamine D<sub>1</sub>/D<sub>2</sub>-receptor agonist apomorphine (43, 44), whereas the GABA<sub>A</sub>-receptor antagonist picrotoxin enhances locomotor stimulant effects of dopamine-receptor agonists (43, 45). Recently, we have found that intra-shell injection of muscimol, but not bicuculline, inhibits the contraversive pivoting induced by a mixture of SKF 38393 and quinpirole injected into the Nacc shell (Fig. 1C) and that this effect of muscimol is antagonized by bicuculline (46), indicating that this inhibitory effect of muscimol on the shell-specific, dopaminergic pivoting is GABA<sub>A</sub>-receptor specific. On the other hand, the shell-specific, dopaminergic pivoting is also inhibited by baclofen, but this effect of baclofen is not prevented by 2-hydroxysaclofen (46), suggesting that the dopamine D<sub>1</sub>/D<sub>2</sub> receptor-mediated pivoting is not mediated by GABA<sub>B</sub> receptors. Since stimulation of GABA<sub>A</sub> receptors inhibits the dopamine release in the Nacc (42), the effect of muscimol is elicited via inhibition of endogenous dopamine release, which causes insufficient stimulation of dopamine D<sub>1</sub>/D<sub>2</sub> receptors. Previously it has been shown that stimulation of D<sub>3</sub> receptors (one of the D<sub>2</sub>-like receptors) reduces GABA<sub>A</sub>-receptor current through a phospho-dependent endocytosis mechanism in the Nacc (47). If the reduction of GABA<sub>A</sub>-receptor current induced by the stimulation of D<sub>3</sub> receptors contributes to eliciting the shell-specific, dopaminergic pivoting, muscimol might strengthen the GABA<sub>A</sub>-receptor current and inhibit

the shell-specific, dopaminergic pivoting.

In conclusion, GABA<sub>A</sub>, but not GABA<sub>B</sub>, receptors in the Nacc shell exert an inhibitory control upon the shell-specific, dopaminergic pivoting (46).

## 2.3. Acetylcholine

As stated in the Introduction, unilateral activation of acetylcholine receptors in the Nacc shell predominantly elicits contraversive circling, whereas unilateral activation of dopamine D<sub>1</sub>/D<sub>2</sub> receptors in the Nacc shell predominantly elicits contraversive pivoting (15, 16). Thus, these results have indicated that both acetylcholine and dopamine receptors in the Nacc shell play a crucial role in eliciting locomotor activity, but that there are subtle differences.

The available literature reveals that there is an interaction between cholinergic and dopaminergic systems in the Nacc. For instance, stimulation of dopamine D<sub>1</sub> receptors has been found to increase acetylcholine release in the Nacc, whereas stimulation of dopamine D<sub>2</sub> receptors has been found to inhibit acetylcholine release in the Nacc (48, 49). These data suggest that dopamine D<sub>1</sub> receptors exert an excitatory control on the cholinergic neurons in contrast to dopamine D<sub>2</sub> receptors that exert an inhibitory control on these neurons. On the other hand, a muscarinic acetylcholine-receptor antagonist has been found to increase dopamine release in the Nacc, whereas a muscarinic acetylcholine-receptor agonist has been found to inhibit the increase of dopamine release induced by amphetamine (50) or by electric stimulation (51), suggesting that muscarinic acetylcholine receptors exert an inhibitory control on the dopaminergic neurons in the Nacc. Moreover, a nicotinic acetylcholine-receptor agonist has been found to increase dopamine release in the Nacc (52), suggesting that nicotinic acetylcholine receptors exert an excitatory control on the dopaminergic neurons.

In line with the above-mentioned neurochemical findings, it has been found that the nicotinic acetylcholine-receptor antagonist mecamylamine significantly suppresses the contraversive pivoting induced by the mixture of SKF 38393 and quinpirole injected into the Nacc shell (Fig. 1D) (29), suggesting that accumbal, nicotinic acetylcholine receptors indeed exert an excitatory control on shell-specific, dopaminergic-dependent behavior as suggested by the above-mentioned neurochemical studies. Since nicotinic acetylcholine receptors are both presynaptically and postsynaptically localized in the Nacc (53), it cannot be excluded that the effect of mecamylamine is just mediated via an enhanced dopamine release. Recently, nicotinic acetylcholine receptors in the Nacc shell have been reported to form complexes with presynaptic dopamine D<sub>2</sub> receptors; stimulation of these nicotinic




receptors has been found to increase dopamine level (54). In contrast, postsynaptic, nicotinic acetylcholine receptors are known to facilitate the GABA-mediated inhibition of the output neurons from the Nacc (53). There is evidence that the contraversive pivoting induced by the stimulation of dopamine D<sub>1</sub>/D<sub>2</sub> receptors in the Nacc shell is funneled via GABAergic neurons innervating the SNc (15, 18, 21) and that GABAergic neurons from the Nacc to the SNc are inhibited by the stimulation of dopamine D<sub>1</sub>/D<sub>2</sub> receptors in the Nacc shell (15). On the basis of these data, it has been suggested that the mecamylamine's inhibitory effects on dopaminergic function are at least partly induced by disinhibition of the GABAergic output neurons from the Nacc shell (53, 55).

The muscarinic acetylcholine-receptor antagonist methylscopolamine significantly increases the contraversive pivoting induced by a mixture of SKF 38393 and quinpirole injected into the Nacc shell (Fig. 2A) (55), suggesting that the muscarinic acetylcholine receptors indeed exert an inhibitory control on the shell-specific, dopamine-dependent behavior as suggested by the above-mentioned neurochemical studies. Given the findings that a muscarinic acetylcholine-receptor antagonist increases dopamine release in the Nacc (50) and muscarinic acetylcholine receptors inhibit dopamine D<sub>1</sub>-receptor signaling in Nacc (56), it has been hypothesized that methylscopolamine potentiates the dopaminergic pivoting by 1) a methylscopolamine-mediated enhancement of dopamine release in the Nacc shell, and by 2) a methylscopolamine-mediated disinhibition of dopamine D<sub>1</sub>-receptor signaling in the Nacc shell (50, 55, 56).

In contrast to the cholinergic influence on the shell-specific, dopamine-dependent behavior, dopamine D<sub>1</sub>/D<sub>2</sub> receptors appear to exert no influence on accumbal, acetylcholine-dependent behavior. Thus, it has been found that inhibition of dopamine D<sub>1</sub> and/or D<sub>2</sub> receptors does not affect the contraversive circling elicited by unilateral administration of the acetylcholine-receptor agonist carbachol into the Nacc shell (55). To what extent these data really indicate that there is no influence of dopamine receptors on the function mediated by cholinergic receptors remains open for discussion because the above-mentioned neurochemical studies have revealed that dopamine D<sub>1</sub> and D<sub>2</sub> receptors exert an excitatory and inhibitory control on the cholinergic neurons, respectively.

The available information has led to the conclusion that stimulation of nicotinic acetylcholine receptors in the Nacc shell is required for the shell-specific, dopamine-dependent behavior and that, in contrast, muscarinic acetylcholine receptors in the Nacc shell inhibit the production of the shell-specific, dopamine-dependent be-



**Fig. 2.** Potentiation of contraversive pivoting induced by a mixture of SKF 38393 and quinpirole injected into the nucleus accumbens shell by intra-shell injections of methylscopolamine (A), isoprenaline (B), orexin A (C), orexin B (D), and somatostatin (E). Methylscopolamine (1 and 2.5 µg), isoprenaline (0.5 and 1 µg), orexin A (1 and 2 µg), orexin B (0.5 and 1 µg), and somatostatin (0.5 and 1 µg) were injected into the nucleus accumbens shell as a cocktail with a mixture of SKF 38393 (1 µg) and quinpirole (10 µg). The contraversive pivoting was measured soon after the drug injections. \* $P < 0.05$ , \*\* $P < 0.01$  vs. SKF 38393 + quinpirole group.

havior (55).

## 2.4. Noradrenaline

The Nacc shell receives noradrenergic projections mainly from the nucleus tractus solitaries and the locus coeruleus (57). Previous studies have revealed that noradrenaline can functionally influence dopamine in the Nacc (58; for review: 59, 60). For example, stimulation of  $\alpha$ -adrenoceptors has been found to decrease dopamine release in the Nacc, whereas stimulation of

$\beta$ -adrenoceptors has been found to increase dopamine release in the Nacc (58, 61). Thus, these results suggest that  $\alpha$ - and  $\beta$ -adrenoceptors exert an inhibitory and an excitatory control on dopamine release in the Nacc, respectively.

To determine the functional interaction between adrenergic and dopaminergic receptors in the Nacc shell, we have analyzed the role of  $\alpha$ - and  $\beta$ -adrenoceptors in the Nacc shell in rat turning behavior. The local administration of neither  $\alpha$ - nor  $\beta$ -adrenoceptor agonists and antagonists elicits turning behavior (62). As described above, although a  $\beta$ -adrenoceptor agonist increases dopamine release in the Nacc (58, 61), it cannot elicit turning behavior. The following possibilities may explain this discrepancy. One is that simultaneous stimulation of both dopamine D<sub>1</sub> and D<sub>2</sub> receptors in the Nacc shell elicits contraversive pivoting, but that stimulation of only one type of dopamine receptors does not elicit turning behavior (17). If increased dopamine stimulates either D<sub>1</sub> or D<sub>2</sub> receptors, it cannot elicit turning behavior. Another possibility is the localization of receptors in the Nacc shell. We have already reported that there are at least two distinct pathways involved in the production of pivoting, namely the shell-ventral pallidum pathway that mediates locomotor activity and the accumbens–nigra–striatum pathway that mediates asymmetry (15, 17, 21). If other neurotransmitters affect just one dopaminergic pathway, they cannot elicit turning behavior.

Unilateral injection of the  $\alpha$ -adrenoceptor agonist phenylephrine into the Nacc shell dose-dependently inhibited contraversive pivoting induced by a mixture of SKF 38393 and quinpirole into the Nacc shell (Fig. 1E) (62). This effect of phenylephrine was antagonized by phentolamine, which alone did not affect the dopaminergic pivoting. Therefore, it can be concluded that the inhibitory effect of phenylephrine on the dopaminergic pivoting is  $\alpha$ -adrenoceptor-specific within the Nacc shell. The fact that  $\alpha$ -adrenoceptors inhibit dopaminergic pivoting might be explained by presynaptic and/or post-synaptic mechanisms. As mentioned above, stimulation of  $\alpha$ -adrenoceptors inhibits dopamine release in the Nacc (58, 60). This outcome implies that stimulation of accumbal dopamine D<sub>1</sub>/D<sub>2</sub> receptors by means of SKF 38393 and quinpirole elicits pivoting only when the baseline tonus at one or both subtypes of these dopamine receptors is already high. Since stimulation of dopamine D<sub>1</sub> receptors by means of endogenous dopamine has been found to be necessary for the display of behavior elicited by stimulation of dopamine D<sub>2</sub> receptors (63), it is thought that the baseline tonus is high, especially at the level of the dopamine D<sub>1</sub> receptors. Accordingly, it is attractive to postulate that accumbal  $\alpha$ -adrenoceptors control at least accumbal dopamine D<sub>1</sub> receptors.

We have shown that the  $\beta$ -adrenoceptor agonist isoprenaline injected into the Nacc shell dose-dependently potentiated the contraversive pivoting induced by a mixture of SKF 38393 and quinpirole injected into the Nacc shell (Fig. 2B) (62); the effect of isoprenaline was antagonized by a dose of propranolol that *per se* had no effect on this pivoting, indicating that the potentiating effect of isoprenaline is  $\beta$ -adrenoceptor-specific (62). These data agree with the findings from neurochemical studies showing that stimulation of  $\beta$ -adrenoceptors increases dopamine release in the Nacc shell (62). All these data indicate that  $\alpha$ -adrenoceptors in the Nacc shell exert an inhibitory effect on the shell-specific, dopaminergic pivoting in contrast to  $\beta$ -adrenoceptors in the Nacc shell, which exert a stimulatory effect on shell-specific dopaminergic pivoting (for references, see above).

## 2.5. Serotonin

Serotonin (5-HT) receptors are widely distributed in the brain (64). There are many reports indicating that the serotonergic system modulates dopamine function in the brain. However, most reports focused on the interaction between 5-HT and dopamine in the VTA, since serotonin neurons project from the dorsal and medial raphe nuclei to the VTA (65). In fact, injection of 5-HT into the VTA increases dopamine level in the Nacc (66). The interaction between 5-HT and dopamine in the Nacc has also been reported. The 5-HT<sub>1B</sub>-receptor agonist increases dopamine level in the Nacc (67). Moreover, 5-HT<sub>2C</sub>-receptor antagonists inhibit cocaine-induced dopamine release in the Nacc (68). In the behavioral studies, the injection of 5-HT<sub>2C</sub>-receptor agonist potentiates cocaine-induced behavior (69). More specifically, injection of 5-HT<sub>1B</sub>-receptor agonist and antagonist into the Nacc shell potentiates and inhibits cocaine-induced locomotor activity, respectively, whereas the same drug injected into the Nacc core is without effect (70). Since the reports examining the role of the serotonergic system in the Nacc shell and core are limited and there are many subtypes of 5-HT receptors, further studies are needed to clarify the role of the serotonergic system in modulating accumbal dopamine functions.

## 2.6. Opioid

Opioid receptors (i.e.,  $\mu$ -,  $\delta$ -, and  $\kappa$ -opioid receptors) have been reported to regulate mesolimbic dopaminergic neuronal activities. For example, activation of  $\mu$ -opioid receptors by intra-cerebral administration of the agonists fentanyl and [D-Ala<sup>2</sup>, N-Me-Phe<sup>4</sup>, Gly<sup>5</sup>-ol]-enkephalin (DAMGO) (VTA: 71, Nacc: 72) and  $\delta$ -opioid receptors by means of the agonists deltorphin II and [D-Pen<sup>2,5</sup>]-enkephalin (DPDPE) given into the Nacc (72, 73) enhances extracellular dopamine concentration in the Nacc,

whereas activation of  $\kappa$ -opioid receptors by means of the agonist U69593 given into the Nacc decreases dopamine concentration in the Nacc (74). The involved receptors are thought to be located in the VTA (for  $\mu$ -opioid receptors) and the Nacc (for  $\mu$ -,  $\delta$ -, and  $\kappa$ -opioid receptors).

Stimulation of  $\mu$ -opioid receptors in the VTA has been found to increase locomotor activity (75, 76). This effect is considered to be dopamine-dependent because it does not occur in 6-hydroxydopamine (6-OHDA)-lesioned animals (76). In contrast, hyperlocomotion induced by  $\mu$ -opioid receptor stimulation in the Nacc is unaffected by dopamine depletion (75, 77). A similar hyperlocomotion induced by  $\delta$ -opioid receptor stimulation in the Nacc is also unaffected by dopamine depletion (73, 77). Therefore, the enhanced locomotor activity induced by injections of  $\mu$ - and  $\delta$ -opioid receptor agonists into the Nacc is thought to be dopamine-independent. However, because 6-OHDA-lesioned animals were used in the latter studies, compensatory changes in the Nacc may have affected the results. This probably explains why these behavioral results are contradictory to the results of the above-mentioned neurochemical studies.

We have examined the role of opioid receptors in the Nacc shell in eliciting dopamine-dependent behavior. Unilateral injections of DAMGO and deltorphin II, but not DPDPE, into the Nacc shell significantly elicited contraversive pivoting (78). The pivoting induced by DAMGO and deltorphin II was receptor-specific because each behavioral response was significantly inhibited by the  $\mu$ -opioid receptor antagonist D-Phe-Cys-Tyr-D-Trp-Orn-Thr-Phe-Thr-NH<sub>2</sub> (CTOP) and the  $\delta$ -opioid receptor antagonist naltriben (NTB), respectively (78). The phenotype of this contraversive pivoting was similar to that elicited by unilateral injection of a mixture of SKF 38393 and quinpirole into the Nacc shell (see also 15, 16, 21). In fact, the contraversive pivoting elicited by stimulation of  $\mu$ - or  $\delta$ -opioid receptors in the Nacc shell was significantly inhibited by *cis*-(Z)-flupenthixol injected into the Nacc shell (78). These results allow the conclusion that the  $\mu$ - and  $\delta$ -opioid receptor-mediated effects require an intact dopaminergic system. This fits in with our earlier published data that local applications of these opioid receptor agonists (i.e., DAMGO and deltorphin II) enhance extracellular levels of dopamine in the Nacc (72). Thus, it is likely that both  $\mu$ - and  $\delta$ -opioid receptors are presynaptically localized on dopaminergic neurons that mediate the dopaminergic pivoting.

Despite the fact that stimulation of  $\delta$ -opioid receptors, like stimulation of  $\mu$ - and  $\delta$ -opioid receptors, enhances the release of accumbal dopamine (72), such stimulation did not elicit pivoting (78). A possible explanation for this finding is that the  $\delta$ -opioid receptors, in contrast to the  $\mu$ - and  $\delta$ -opioid receptors, are localized on dopamin-

ergic terminals that are not involved in the production of pivoting. Indeed, there are at least two distinct neuronal pathways involved in the production of pivoting, namely the shell-ventral pallidum pathway that mediates locomotor activity and the accumbens-nigra-striatum pathway that mediates asymmetry (17; see also 16, 21). In view of these considerations, it has been hypothesized that a)  $\mu$ - and  $\delta$ -opioid receptors are presynaptically localized on dopaminergic neurons that impinge on both the shell-ventral pallidum pathway and the accumbens-nigra-striatum pathway, and that b)  $\delta$ -opioid receptors are at best presynaptically localized on dopaminergic neurons that impinge just on the accumbens-nigra-striatum pathway (72, 73, 78). An alternative explanation for the ineffectiveness of the  $\delta$ -opioid receptor agonist might be that the DPDPE-induced increase of extracellular dopamine is due not to stimulation of  $\delta$ -opioid receptors, but to stimulation of opioid-independent mechanisms.

It has been found that contraversive pivoting induced by a mixture of SKF 38393 and quinpirole into the Nacc shell is also inhibited by NTB (Fig. 1F), but not by CTOP or the  $\delta$ -opioid receptor antagonist (*E*)-7-benzyliden-naltrexone (BNTX) (78). These results therefore suggest that  $\delta$ -, but not  $\mu$ - and  $\delta$ -, opioid receptors interact with postsynaptically localized dopamine D<sub>1</sub>/D<sub>2</sub>-receptor mechanisms involved in the production of pivoting.

These results demonstrate that unilateral stimulations of  $\mu$ - or  $\delta$ -opioid receptors, but not  $\delta$ -opioid receptors, in the Nacc shell elicit contraversive pivoting that requires intact dopamine D<sub>1</sub>/D<sub>2</sub>-receptor mechanisms in the Nacc shell. Moreover, it has been shown that the shell-specific, dopaminergic pivoting is modulated by  $\delta$ -, but not  $\mu$ - and  $\delta$ -, opioid receptors.

## 2.7. Orexin

Orexins (orexin A and B) are peptides, which were originally isolated from rat hypothalamus (79). Because the hypothalamus is the regulatory center for autonomic and endocrine homeostatic systems, orexins are thought to regulate such systems including feeding behavior, arousal activity, and arterial pressure (79–81). Orexin neurons and orexin receptors are not only present in the hypothalamus, but also extensively distributed across various brain areas (82, 83). This may imply that these peptides play an important role in several physiological functions.

Previously, it has been reported that orexin neurons originating in the lateral hypothalamus impinge upon dopaminergic cells in the VTA (84–86). Orexins injected into the VTA activate dopaminergic neurons (86). These results indicate that stimulation of orexin receptors in the VTA activates dopamine neurons and, accordingly,



increases dopamine release in the Nacc (86), and, subsequently, increases locomotor activity.

Since the Nacc has been found to contain mRNA for both orexin 1 receptors and orexin 2 receptors (83), we examined the role of orexin receptors in dopamine-dependent behavior. Orexin A injected into the Nacc shell significantly potentiated the contraversive pivoting induced by a mixture of SKF 38393 and quinpirole injected unilaterally into the Nacc shell (Fig. 2C) (87). The potentiating effect of orexin A was not antagonized by the orexin 1-receptor antagonist SB 334867, which alone did not significantly modify the contraversive pivoting induced by the mixture of SKF 38393 and quinpirole injected unilaterally into the Nacc shell (87). Our result indicates that the effect of orexin A was not mediated via orexin 1 receptors in the Nacc shell. Since orexin A is equipotent at orexin 1 and 2 receptors, we have examined the role of orexin 2 receptors in dopaminergic pivoting. The orexin 2 receptor-selective agonist orexin B injected into the Nacc shell significantly potentiated the contraversive pivoting induced by the mixture of SKF 38393 and quinpirole injected unilaterally into the Nacc shell (Fig. 2D) (87). This result shows that orexin 2 receptors play an important, modifying role in eliciting the shell-specific, dopaminergic pivoting. This conclusion is supported by the recent report that orexin B and dopamine synergistically increases the firing rate of neurons in the Nacc shell (88).

Previous reports have indicated that the cell bodies of dopaminergic neurons in the VTA are innervated by orexin neurons originating in the lateral hypothalamus and that stimulation of orexin receptors in the VTA activates dopamine neurons (84–86). Since the injection of orexin A into the VTA is reported to be more effective in eliciting hyperlocomotion than that of orexin B (85, 86), it has been suggested that orexin 1 receptors, but not orexin 2 receptors, in the VTA stimulate dopaminergic neurons. Our results demonstrate that orexin 2 receptors in the Nacc shell, which is the terminal area of dopamine neurons originating in the VTA, also modulate the function mediated by accumbal dopamine. These data are in line with the previously reported finding that the expression of orexin 2-receptor mRNA in the Nacc is greater than that of orexin 1-receptor mRNA in the Nacc (83). Thus, different subtypes of orexin receptors, that is, orexin 1 and 2 receptors, modulate dopaminergic function in the VTA and the Nacc, respectively.

## 2.8. Somatostatin

Somatostatin is a tetradecapeptide that was originally isolated from mammalian hypothalamus and characterized as a potent physiological inhibitor of growth hormone secretion from the anterior pituitary (89). Soma-

tostatin and its receptors are not only present in the hypothalamus, but also widely distributed across the central nervous system (90, 91). Since the striatum and Nacc are reported to contain both somatostatin and its receptors (90, 92), it is possible that somatostatin regulates dopaminergic function in these brain areas.

Many neurochemical and behavioral studies on the functional role of somatostatin in the striatum have been reported. In neurochemical studies, systemic or intra-striatal injection of somatostatin or a somatostatin-receptor agonist increases dopamine release in the striatum (93). These results indicate that somatostatin positively regulates dopamine release in the striatum. Conversely, dopamine also regulates somatostatin level in the striatum (94, 95). For example, dopamine-receptor antagonists such as haloperidol reduce somatostatin-like immunoreactivity in the striatum (94). Moreover, dopamine D<sub>1</sub>- and D<sub>2</sub>-receptor agonists potentiate the effects of somatostatin on somatostatin-receptor binding, inhibition of adenylyl cyclase activity, and accumulation of inositol 1,4,5-trisphosphate (95). In behavioral studies, somatostatin in the striatum has also been found to play a regulatory role in dopamine-dependent behavior. Intra-striatal injection of cysteamine, which selectively reduces somatostatin levels, blocks apomorphine-induced stereotypy (96). These neurochemical and behavioral findings show that somatostatin in the striatum plays an important role in dopaminergic functions.

In contrast to the striatum, the role of somatostatin in the Nacc has received less attention. Neurochemical studies have indicated that the perfusion of somatostatin into the Nacc increases dopamine release in the same area (97). To determine the function of accumbal somatostatin on dopamine-dependent behavior, we examined the role of somatostatin on dopamine-dependent behavior. Somatostatin significantly potentiated the contraversive pivoting induced by the mixture of SKF 38393 and quinpirole unilaterally injected into the Nacc shell (Fig. 2E) (98). This potentiating effect of somatostatin was significantly antagonized by the somatostatin-receptor antagonist cyclosomatostatin, which alone did not significantly modify the contraversive pivoting induced by the mixture of SKF 38393 and quinpirole injected unilaterally into the Nacc shell (99). Somatostatin has been reported to increase dopamine release in the Nacc (97). Moreover, in the striatum, somatostatin increases dopamine release via the glutamatergic system, especially via AMPA and NMDA receptors (93). We have shown that the shell-specific, dopaminergic pivoting is mediated by both AMPA and NMDA receptors in the same area (see above and ref. 35). Therefore, one can speculate that somatostatin increases dopamine release via the glutamatergic system, including AMPA and NMDA receptors,

in the Nacc shell. Another possibility is that somatostatin increases dopamine release caused by inhibiting GABA release, since somatostatin has been found to regulate GABA release presynaptically in the striatum (99). Thus, the potentiating effect of somatostatin on the contraversive pivoting induced by dopamine D<sub>1</sub>/D<sub>2</sub>-receptor stimulation in the Nacc shell may be due to additional stimulation of dopamine D<sub>1</sub> and D<sub>2</sub> receptors caused by increased dopamine. It is also possible that somatostatin regulates dopamine-receptor functions postsynaptically in the potentiating effect of somatostatin because dopamine and somatostatin-receptor activation may work in a co-operative manner on postsynaptic sites, possibly via second messenger pathways or common targets (100).

### 3. Conclusion

The present study focused on the modulation of the shell-specific, dopaminergic pivoting by glutamate, GABA, noradrenaline, acetylcholine, opioid, orexin, and somatostatin in the Nacc shell: these modulations vary across the different neurotransmitters, neuropeptides, and their corresponding receptors (Table 1). It has been shown that the dopaminergic pivoting is funneled from the Nacc shell via the SNc, the ventrolateral striatum, and the substantia nigra pars reticulata towards the peduncu-

lopontine tegmental nucleus (15, 18, 21). Thus, this pathway is likely to be controlled by these neurotransmitters in the Nacc shell, implying that all functions mediated via this pathway are sensitive to modulation by compounds that selectively interact with one or more of these neuroactive compounds. As mentioned above, the shell-specific, dopaminergic pivoting is just used as a tool to trace the nature of the interactions of dopamine with other neuroactive compounds and *vice versa* as well as a tool to trace the pathways involved in the transmission of accumbal dopamine-dependent information across the brain: it has no face validity with any of the known functions of accumbal dopamine. Given the well-known therapeutic effects of drugs that enhance accumbal dopamine, in patients suffering from depression, ADHD or Parkinson's disease, agents that directly or indirectly stimulate the activity at the level of the receptors shown in Table 1, are predicted to have a therapeutic efficacy as well. On the other hand, it is predicted that agents that directly or indirectly inhibit the activity at the level of the receptors shown in Table 1, are therapeutically effective in patients suffering from schizophrenia, because drugs that reduce the accumbal dopaminergic activity in such patients are therapeutically effective. The putative advantage of using drugs that indirectly alter the activity at the level of the accumbal dopamine receptors is that

**Table 1.** Summary of the role of each neuroactive compound on accumbens-specific behavior

		Production of turning behavior	Modulation of dopamine-dependent turning behavior
Glutamate receptors	NMDA*	—	↑
	AMPA*	↑	↑
GABA receptors	GABA <sub>A</sub>	—	↓
	GABA <sub>B</sub>	—	—
Acetylcholine receptors	muscarine	↑	↓
	nicotine	↑	↑
Adrenaline receptors	$\alpha$	—	↓
	$\beta$	—	↑
Serotonin receptors	5-HT <sub>1B</sub>	—	↑
	5-HT <sub>2C</sub>	—	↑
Opioid receptors	$\mu$	↑	—
	$\delta_1$	—	—
	$\delta_2$	↑	↑
Orexin receptors	orexin 1	—	—
	orexin 2	—	↑
Somatostatin receptors		—	↑

\*Stimulation indirectly affects accumbal dopamine-dependent pivoting. Up arrow: increases behavior, down arrow: decreases behavior.

such drugs do not alter the sensitivity of the involved dopamine receptors and, therefore, result in less unwanted side-effects than drugs that directly affect the dopamine receptors.

In conclusion, since both accumbal dopamine, as well as the above-mentioned pathway, play a crucial role in a great variety of neurological and psychiatric diseases such as depression, drug addiction, ADHD, schizophrenia, and Parkinson's disease, it is suggested that drugs that selectively interfere with one or more of these non-dopaminergic compounds, are promising candidates for treating these diseases.

## Acknowledgments

This study was partly supported by a grant for promotion of multi-disciplinary research projects from the Ministry of Education, Culture, Sports, Science, and Technology (N.K., A.R.C.); the Promotion and Mutual Aid Corporation for Private Schools in Japan (H.I., N.K.); and Grant-in-Aid for Scientific Research (23792133 to H.I.) from Japan Society for Promotion of Science.

## References

- Kienast T, Heinz A. Dopamine and the diseased brain. *CNS Neurol Disord Drug Targets*. 2006;5:109–131.
- Nestler EJ, Carlezon WA Jr. The mesolimbic dopamine reward circuit in depression. *Biol Psychiat*. 2006;59:1151–1159.
- Di Chiara G, Bassareo V, Fenu S, de Luca MA, Spina L, Cadoni C, et al. Dopamine and drug addiction: the nucleus accumbens shell connection. *Neuropharmacology*. 2004;47 Suppl 1:227–241.
- Heimer L. A new anatomical framework for neuropsychiatric disorders and drug abuse. *Am J Psychiat*. 2003;160:1726–1739.
- Yamada K, Nagai T, Nabeshima T. Drug dependence, synaptic plasticity, and tissue plasminogen activator. *J Pharmacol Sci*. 2005;97:157–161.
- Levy F. Synaptic gating and ADHD: a biological theory of comorbidity of ADHD and anxiety. *Neuropsychopharmacology*. 2004;29:1589–1596.
- Sounga-Barke EJS. Causal models of attention-deficit/hyperactivity disorder: from common simple deficits to multiple developmental pathways. *Biol Psychiat*. 2005;57:1231–1238.
- Grace AA. Gating of information flow within the limbic system and the pathophysiology of schizophrenia. *Brain Res Rev*. 2000;31:330–341.
- Pani L. Clinical implications of dopamine research in schizophrenia. *Curr Med Res Opin*. 2002;18 Suppl 3:s3–s7.
- Olanow CW, Obeso JA, Stocchi F. Continuous dopamine-receptor treatment of Parkinson's disease: rationale and clinical implications. *Lancet Neurol*. 2006;5:677–687.
- Bardo MT, Hammer RP Jr. Autoradiographic localization of dopamine D<sub>1</sub> and D<sub>2</sub> receptors in rat nucleus accumbens: resistance to differential conditions. *Neuroscience*. 1991;45:281–290.
- Deutch AY, Cameron DS. Pharmacological characterization of dopamine systems in the nucleus accumbens core and shell. *Neuroscience*. 1992;46:49–56.
- Voorn P, Jorritsma-Byham B, van Dijk C, Buijs RM. The dopaminergic innervation of the ventral striatum in the rat: a light- and electron-microscopical study with antibodies against dopamine. *J Comp Neurol*. 1986;251:84–99.
- Zahm DS, Heimer L. Specificity in the efferent projections of the nucleus accumbens in the rat: comparison of the rostral pole projection patterns with those of core and shell. *J Comp Neurol*. 1993;327:220–232.
- Kitamura M, Koshikawa N, Yoneshige N, Cools AR. Behavioural and neurochemical effects of cholinergic and dopaminergic agonists administered into the accumbal core and shell in rats. *Neuropharmacology*. 1999;38:1397–1407.
- Koshikawa N, Kitamura M, Kobayashi M, Cools AR. Contralateral turning elicited by unilateral stimulation of dopamine D<sub>2</sub> and D<sub>1</sub> receptors in the nucleus accumbens of rats is due to stimulation of these receptors in the shell, but not the core, of this nucleus. *Psychopharmacology*. 1996;126:185–190.
- Koshikawa N. Role of the nucleus accumbens and the striatum in the production of turning behaviour in intact rats. *Rev Neurosci*. 1994;5:331–346.
- Ikeda H, Akiyama G, Matsuzaki S, Sato M, Koshikawa N, Cools AR. GABA<sub>A</sub> receptors in the pedunculopontine tegmental nucleus play a crucial role in rat shell-specific dopamine-mediated, but not shell-specific acetylcholine-mediated, turning behaviour. *Neuroscience*. 2004;125:553–562.
- Ikeda H, Kotani A, Koshikawa N, Cools AR. Differential role of GABA<sub>A</sub> and GABA<sub>B</sub> receptors in two distinct output stations of the rat striatum: studies on the substantia nigra pars reticulata and the globus pallidus. *Neuroscience*. 2010;167:31–39.
- Ikeda H, Kotani A, Lee J, Koshikawa N, Cools AR. GABA<sub>A</sub> receptors in the mediodorsal thalamus play a crucial role in rat shell-specific acetylcholine-mediated, but not dopamine-mediated, turning behavior. *Neuroscience*. 2009;159:1200–1207.
- Kitamura M, Ikeda H, Koshikawa N, Cools AR. GABA<sub>A</sub> agents injected into the ventral pallidum differentially affect dopaminergic pivoting and cholinergic circling elicited from the shell of the nucleus accumbens. *Neuroscience*. 2001;104:117–127.
- Maggio R, Aloisi G, Silvano E, Rossi M, Millan MJ. Heterodimerization of dopamine receptors: new insights into functional and therapeutic significance. *Parkinsonism Relat Disord*. 2009;15 Suppl 4:S2–S7.
- Missale C, Fiorentini C, Collo G, Spano P. The neurobiology of dopamine receptors: evolution from the dual concept to heterodimer complexes. *J Recept Signal Transduct Res*. 2010;30:347–354.
- Kearn CS, Blake-Palmer K, Daniel E, Mackie K, Glass M. Concurrent stimulation of cannabinoid CB1 and dopamine D2 receptors enhances heterodimer formation: a mechanism for receptor cross-talk? *Mol Pharmacol*. 2005;67:1697–1704.
- Kelly AE, Domesick VB. The distribution of projection from the hippocampal formation to the nucleus accumbens in the rat: an anterograde and retrograde horseradish peroxidase study. *Neuroscience*. 1982;7:2321–2335.
- Robinson TG, Beart PM. Excitant amino acid projections from rat amygdala and thalamus to the nucleus accumbens. *Brain Res Bull*. 1988;20:467–471.
- Walaas I. Biochemical evidence for overlapping neocortical and allocortical glutamate projections to the nucleus accumbens and rostral caudatoputamen in the rat brain. *Neuroscience*. 1981;6:399–405.
- Sesack SR, Pickel VM. In the rat medial nucleus accumbens,

- hippocampal and catecholaminergic terminals converge on spiny neurons and are in apposition to each other. *Brain Res.* 1990; 527:266–279.
- 29 Sesack SR, Pickel VM. Prefrontal cortical efferents in the rat synapse on unlabeled neuronal targets of catecholamine terminals in the nucleus accumbens septi and on dopamine neurons in the ventral tegmental area. *J Comp Neurol.* 1992;320:145–160.
  - 30 Young AB, Fagg GE. Excitatory amino acid receptors in the brain: membrane binding and receptor autoradiographic approaches. 1990;11:126–133.
  - 31 Imperato A, Honoré T, Jensen LH. Dopamine release in the nucleus caudatus and in the nucleus accumbens is under glutamatergic control through non-NMDA receptors: a study in free-moving rats. *Brain Res.* 1990;530:223–228.
  - 32 Dalia A, Uretsky NJ, Wallace LJ. Dopaminergic agonists administered into the nucleus accumbens: Effects on extracellular glutamate and on locomotor activity. *Brain Res.* 1998;788:111–117.
  - 33 Boldry RC, Uretsky NJ. The importance of dopaminergic neurotransmission in the hypermotility response produced by the administration of N-methyl-D-aspartic acid into the nucleus accumbens. *Neuropharmacology.* 1988;27:569–577.
  - 34 Boldry RC, Willins DL, Wallace LJ, Uretsky NJ. The role of endogenous dopamine in the hypermotility response to intra-accumbens AMPA. *Brain Res.* 1991;559:100–108.
  - 35 Ikeda H, Akiyama G, Fujii Y, Minowa R, Koshikawa N, Cools AR. Role of AMPA and NMDA receptors in the nucleus accumbens shell in turning behavior of rats: interaction with dopamine receptors. *Neuropharmacology.* 2003;44:81–87.
  - 36 Taber MT, Baker GB, Fibiger HC. Glutamate receptor agonists decrease extracellular dopamine in the rat nucleus accumbens *in vivo*. *Synapse.* 1996;24:165–172.
  - 37 Kaddis FG, Wallace LJ, Uretsky NJ. AMPA/kainate antagonists in the nucleus accumbens inhibit locomotor stimulatory response to cocaine and dopamine agonists. *Pharmacol Biochem Behav.* 1993;46:703–708.
  - 38 Pulvirenti L, Swerdlow NR, Koob GF. Nucleus accumbens NMDA antagonist decreases locomotor activity induced by cocaine, heroin or accumbens dopamine, but not caffeine. *Pharmacol Biochem Behav.* 1991;40:841–845.
  - 39 Hara Y, Pickel VM. Overlapping intracellular and differential synaptic distributions of dopamine D1 and glutamate N-methyl-D-aspartate receptors in rat nucleus accumbens. *J Comp Neurol.* 2005;492:442–455.
  - 40 Carr KD, Chau LS, de Vaca C, Gustafson K, Stouffer M, Tukey DS, et al. AMPA receptor subunit GluR1 downstream of D-1 dopamine receptor stimulation in nucleus accumbens shell mediates increased drug reward magnitude in food-restricted rats. *Neuroscience.* 2010;165:1074–1086.
  - 41 Matsumoto RR. GABA receptors: are cellular differences reflected in function? *Brain Res Rev.* 1989;14:203–225.
  - 42 Rahman S, McBride WJ. Involvement of GABA and cholinergic receptors in the nucleus accumbens on feedback control of somatodendritic dopamine release in the ventral tegmental area. *J Neurochem.* 2002;80:646–654.
  - 43 Pycck CJ, Horton RW. Dopamine-dependent hyperactivity in the rat following manipulation of GABA mechanisms in the region of the nucleus accumbens. *J Neural Trans.* 1979;45:17–33.
  - 44 Scheel-Krüger J, Cools AR, Van Wel PM. Muscimol a GABA-agonist injected into the nucleus accumbens increases apomorphine stereotypy and decreases the motility. *Life Sci.* 1977;21:1697–1702.
  - 45 Wong LS, Eshel G, Dreher J, Org J, Jackson DM. Role of dopamine and GABA in the control of motor activity elicited from the rat nucleus accumbens. *Pharmacol Biochem Behav.* 1991; 38:829–835.
  - 46 Akiyama G, Ikeda H, Matsuzaki S, Sato M, Moribe S, Koshikawa N, et al. GABA<sub>A</sub> and GABA<sub>B</sub> receptors in the nucleus accumbens shell differentially modulate dopamine and acetylcholine receptor-mediated turning behaviour. *Neuropharmacology.* 2004;46: 1082–1088.
  - 47 Chen G, Kittler JT, Moss SJ, Yan Z. Dopamine D<sub>3</sub> receptors regulate GABA<sub>A</sub> receptor function through a phospho-dependent endocytosis mechanism in nucleus accumbens. *J Neurosci.* 2006;26:2513–2521.
  - 48 Consolo S, Caltavuturo C, Colli E, Recchia M, Di Chiara G. Different sensitivity of *in vivo* acetylcholine transmission to D<sub>1</sub> receptor stimulation in shell and core of the nucleus accumbens. *Neuroscience.* 1999;89:1209–1217.
  - 49 Keys AS, Mark GP. D<sub>1</sub> and D<sub>2</sub> dopamine receptor mediation of amphetamine-induced acetylcholine release in nucleus accumbens. *Neuroscience.* 1998;86:521–531.
  - 50 Ichikawa J, Chung YC, Li Z, Dai J, Meltzer HY. Cholinergic modulation of basal and amphetamine-induced dopamine release in rat medial prefrontal cortex and nucleus accumbens. *Brain Res.* 2002;958:176–184.
  - 51 Threlfell S, Clements MA, Khodai T, Pienaar IS, Exley R, Wess J. Striatal muscarinic receptors promote activity dependence of dopamine transmission via distinct receptor subtypes on cholinergic interneurons in ventral versus dorsal striatum. *J Neurosci.* 2010;30:3398–3408.
  - 52 Fu Y, Matta G, Gao W, Sharp BM. Local  $\alpha$ -bungarotoxin-sensitive nicotinic receptors in the nucleus accumbens modulate nicotine-stimulated dopamine secretion *in vivo*. *Neuroscience.* 2000;101:369–375.
  - 53 de Rover M, Lodder JC, Kits KS, Schoffelemeier ANM, Brussaard AB. Cholinergic modulation of nucleus accumbens medium spiny neurones. *Eur J Neurosci.* 2002;16:2279–2290.
  - 54 Quarta D, Ciruela F, Patkar K, Borycz J, Solinas M, Lluís C, et al. Heteromeric nicotinic acetylcholine-dopamine autoreceptor complexes modulate striatal dopamine release. *Neuropsychopharmacology.* 2007;32:35–42.
  - 55 Moribe S, Ikeda H, Sato M, Akiyama G, Matsuzaki S, Hasegawa K, et al. Acetylcholine receptor effects on accumbal shell dopamine-mediated turning behaviour in rats. *Neuropharmacology.* 2005;49:514–524.
  - 56 Onali P, Olanas MC. Muscarinic M<sub>4</sub> receptors inhibition of dopamine D<sub>1</sub>-like receptor signalling in rat nucleus accumbens. *Eur J Pharmacol.* 2002;448:105–111.
  - 57 Delfs JM, Zhu Y, Druhan JP, Aston-Jones GS. Origin of noradrenergic afferents to the shell subregion of the nucleus accumbens: anterograde and retrograde tract-tracing studies in the rat. *Brain Res.* 1998;806:127–140.
  - 58 Cools AR, van den Bos R, Ploeger G, Ellenbroek BA. Gating function of noradrenaline in the ventral striatum: its role in behavioural responses to environmental and pharmacological challenges. In: Willner P, Scheel-Krüger J, editors. *The mesolimbic dopamine system: from motivation to action*. Chichester: John Wiley & Sons Ltd.; 1991. p. 141–173.
  - 59 Verheij MM, Cools AR. Twenty years of dopamine research: individual differences in the response of accumbal dopamine to



- environmental and pharmacological challenges. *Eur J Pharmacol.* 2008;585:228–244.
- 60 Verheij MM, Cools AR. Accumbal noradrenaline that contributes to the alpha-adrenoceptor-mediated release of dopamine from reserpine-sensitive storage vesicles in the nucleus accumbens is derived from alpha-methyl-para-tyrosine-sensitive pools. *J Neural Transm.* 2009;116:389–394.
  - 61 Verheij MM, Cools AR. Mesolimbic alpha-, but not beta-adrenoceptors control the accumbal release of dopamine that is derived from reserpine-sensitive storage vesicles. *Neuroscience.* 2009;162:1163–1173.
  - 62 Ikeda H, Moribe S, Sato M, Kotani A, Koshikawa N, Cools AR. Adrenergic receptors in the nucleus accumbens shell differentially modulate dopamine and acetylcholine receptor-mediated turning behaviour. *Eur J Pharmacol.* 2007;554:175–182.
  - 63 White FJ, Bednars LM, Wachtel SR, Hjorth S, Brooderson RJ. Is stimulation of both D1 and D2 receptors necessary for the expression of dopamine-mediated behaviors? *Pharmacol Biochem Behav.* 1988;30:189–193.
  - 64 Barns NM, Sharp T. A review of central 5-HT receptors and their function. *Neuropharmacology.* 1999;38:1083–1152.
  - 65 Moore RY, Halaris AE, Jones BE. Serotonin neurons in the mid-brain raphe: ascending projections. *J Comp Neurol.* 1978;180:417–438.
  - 66 Guan XM, McBride WJ. Serotonin microinfusion into the ventral tegmental area increases accumbens dopamine release. *Brain Res Bull.* 1989;23:541–547.
  - 67 Yan Q-S, Yan S-E. Activation of 5-HT<sub>1B/1D</sub> receptors in the mesolimbic dopamine system increases dopamine release from the nucleus accumbens: a microdialysis study. *Eur J Pharmacol.* 2001;418:55–64.
  - 68 Zaraya AE, McIver G, Valdivia PN, Lominac KD, McCreary AC, Szumlinski KK. Blockade of nucleus accumbens 5-HT<sub>2A</sub> and 5-HT<sub>2C</sub> receptors prevents the expression of cocaine-induced behavioral and neurochemical sensitization in rats. *Psychopharmacology.* 2011;213:321–335.
  - 69 Filip M, Cunningham KA. Serotonin 5-HT<sub>2C</sub> receptors in nucleus accumbens regulate expression of the hyperlocomotive and discriminative stimulus effects of cocaine. *Pharmacol Biochem Behav.* 2002;71:745–756.
  - 70 Przegalinski E, Filip M, Papla I, Czepiel K. Effects of 5-HT<sub>1B</sub> receptor ligands microinjected into the accumbal shell or core on the cocaine-induced locomotor hyperactivity in rats. *J Physiol Pharmacol.* 2002;53:383–394.
  - 71 Spanagel R, Herz A, Shippenberg TS. The effects of opioid peptide on dopamine release in the nucleus accumbens: an in vivo microdialysis study. *J Neurochem.* 1990;55:1734–1740.
  - 72 Yoshida Y, Koide S, Hirose N, Takada K, Tomiyama K, Koshikawa N, et al. Fentanyl increases dopamine release in rat nucleus accumbens: involvement of mesolimbic mu- and delta-2-opioid receptors. *Neuroscience.* 1999;92:1357–1365.
  - 73 Longoni R, Spina L, Mulas A, Carboni E, Garau L, Melchiorri P, et al. (D-Ala<sup>2</sup>)Deltorphin II: D<sub>1</sub>-dependent stereotypies and stimulation of dopamine release in the nucleus accumbens. *J Neurosci.* 1991;11:1565–1576.
  - 74 Spanagel R, Herz A, Shippenberg TS. Opposing tonically active endogenous opioid systems modulate the mesolimbic dopaminergic pathway. *Proc Natl Acad Sci U S A.* 1992;89:2045–2050.
  - 75 Kalivas PW, Widerlöv E, Stanley D, Breese G, Prange AJ Jr. Enkephalin action on the mesolimbic system: a dopamine-dependent and dopamine-independent increase in locomotor activity. *J Pharmacol Exp Ther.* 1983;227:229–237.
  - 76 Stinus L, Koob GF, Ling N, Bloom FE, Moal ML. Locomotor activation induced by infusion of endorphins into the ventral tegmental area: evidence for opiate-dopamine interactions. *Proc Natl Acad Sci U S A.* 1980;77:2323–2327.
  - 77 Churchill L, Kalivas PW. Dopamine depletion produces augmented behavioral responses to a mu-, but not a delta-opioid receptor agonist in the nucleus accumbens: lack of a role for receptor upregulation. *Synapse.* 1992;11:47–57.
  - 78 Matsuzaki S, Ikeda H, Akiyama G, Sato M, Moribe S, Suzuki T, et al. Role of  $\mu$ - and  $\delta$ -opioid receptors in the nucleus accumbens in turning behavior of rats. *Neuropharmacology.* 2004;46:1089–1096.
  - 79 Sakurai T, Amemiya A, Ishii M, Matsuzaki I, Chemelli RM, Tanaka H, et al. Orexins and orexin receptors: a family of hypothalamic neuropeptides and G protein-coupled receptors that regulate feeding behavior. *Cell.* 1998;92:573–585.
  - 80 Sakurai T. The neural circuit of orexin (hypocretin): maintaining sleeping and wakefulness. *Nat Rev Neurosci.* 2007;8:171–181.
  - 81 Samson WK, Gosnell B, Chang J-K, Resch ZT, Murphy TC. Cardiovascular regulatory actions of the hypocretins in brain. *Brain Res.* 1999;831:248–253.
  - 82 Peyron C, Tighe DK, van den Pol AN, de Lecea L, Heller HC, Sutcliffe JG, et al. Neurons containing hypocretin (orexin) project to multiple neuronal systems. *J Neurosci.* 1998;18:9996–10015.
  - 83 Trivedi P, Yu H, MacNeil DJ, Van der Plog LHT, Guan X-M. Distribution of orexin receptor mRNA in the rat brain. *FEBS Lett.* 1998;438:71–75.
  - 84 Fadel J, Deutch Y. Anatomical substrates of orexin-dopamine interactions: lateral hypothalamus projections to the ventral tegmental area. *Neuroscience.* 2002;111:379–387.
  - 85 Nakamura T, Uramura K, Nambu T, Yada T, Goto K, Yanagisawa M, et al. Orexin-induced hyperlocomotion and stereotypy are mediated by the dopaminergic system. *Brain Res.* 2000;873:181–187.
  - 86 Narita M, Nagumo Y, Hashimoto S, Narita M, Khotib J, Miyatake M, et al. Direct involvement of orexinergic systems in the activation of the mesolimbic dopamine pathway and related behaviors induced by morphine. *J Neurosci.* 2006;26:398–405.
  - 87 Kotani A, Ikeda H, Koshikawa N, Cools AR. Role of orexin receptors in the nucleus accumbens in dopamine-dependent turning behavior of rats. *Neuropharmacology.* 2008;54:613–619.
  - 88 Mori K, Kim J, Sasaki K. Electrophysiological effects of orexin-B and dopamine on rat nucleus accumbens shell neurons in vitro. *Peptides.* 2011;32:246–252.
  - 89 Brazeau P, Vale W, Burgus R, Ling M, Butcher J, Rivier J, et al. Hypothalamic polypeptide that inhibits the secretion of immunoreactive pituitary growth hormone. *Science.* 1973;179:77–79.
  - 90 Brownstein M, Arimura A, Sato H, Schally AV, Kizer JS. The regional distribution of somatostatin in the rat brain. *Endocrinology.* 1975;96:1456–1461.
  - 91 Martin JL, Chesselet M-F, Raynor K, Gonzales C, Reisine T. Differential distribution of somatostatin receptor subtypes in rat brain revealed by newly developed somatostatin analogs. *Neuroscience.* 1991;41:581–593.
  - 92 Beal MF, Domesick VB, Martin JB. Regional somatostatin distribution in the rat striatum. *Brain Res.* 1983;278:103–108.
  - 93 Hathway GJ, Emson PC, Humphrey PPA, Kendrick KM. Soma-

- tostatin potently stimulate in vivo striatal dopamine and  $\gamma$ -aminobutyric acid release by a glutamate-dependent action. *J Neurochem.* 1998;70:1740–1749.
- 94 Beal MF, Martin JB. Effects of neuroleptic drugs on brain somatostatin-like-immunoreactivity. *Neurosci Lett.* 1984;47:125–130.
- 95 Izquierdo-Claros RM, Boyano-Adánez MC, Larsson C, Gustavsson L, Arilla E. Acute effects of D<sub>1</sub>- and D<sub>2</sub>-receptor agonist and antagonist drugs on somatostatin binding, inhibition of adenylyl cyclase activity and accumulation of inositol 1,4,5-trisphosphate in the rat striatum. *Mol Brain Res.* 1997;47:99–107.
- 96 Lee N, Radke M, Vincent SR. Intra-cerebral cysteamine infusions attenuate the motor response to dopaminergic agonists. *Behav Brain Res.* 1988;29:179–183.
- 97 Pallis E, Thermos K, Spyraiki C. Chronic desipramine treatment selectively potentiates somatostatin-induced dopamine release in the nucleus accumbens. *Eur J Neurosci.* 2001;14:763–767.
- 98 Ikeda H, Kotani A, Koshikawa N, Cools AR. Somatostatin receptors in the nucleus accumbens modulate dopamine-dependent but not acetylcholine-dependent turning behavior of rats. *Neuroscience.* 2009;159:974–981.
- 99 Lopez-Huerta VG, Tecuapetla F, Guzman JN, Bargas J, Galarraga E. Presynaptic modulation of somatostatin in the neostriatum. *Neurochem Res.* 2008;33:1452–1458.
- 100 Rocheville M, Lange DC, Kumar U, Patel SC, Patel RC, Patel YC. Receptors for dopamine and somatostatin: formation of hetero-oligomers with enhanced functional activity. *Science.* 2000;288:154–157.